

ISSN 2456-4990

J Ayu Med Sci

Quarterly
Journal for
Rapid
Publication
of Researches
in Ayurveda
and Other
Traditional
Medicines



Journal of Ayurveda Medical Sciences

J Ayu Med Sci | 2018 | Vol 3 | Issue 1 (Jan – Mar)

www.jayumedsci.com
©Journal of Ayurveda Medical Sciences

Identification of *Ficus religiosa* by DNA SCAR Marker

Suma Venkatesh Mallya*, Vishwanath Udupi¹, Suchitra Narayana Prabhu¹, Sunil Kumar Koppala Narayana²


Department of Dravyaguna, SDM College of Ayurveda, Kuthpady, ¹Research officer, SDM Centre for Research in Ayurveda & Allied Sciences, Udupi 574118 Karnataka, ²Department of Pharmacognosy, Siddha Central Research Institute, Arumbakkam, Chennai 600106 Tamilnadu, India.

ABSTRACT

Introduction: *Ficus religiosa* Linn. is a popularly used bark drug used in Indian system of medicine, an ingredient of *Panchavalka* (Five bark drugs). Because of morphological similarity adulteration is common with bark where quality therapeutics is always uncertain. DNA markers are reliable in this regard as the genetic composition is unique for each species and is not affected by age, habitat, environmental factors and physiological conditions. Hence DNA SCAR marker development of *F. religiosa* was carried out. **Methods:** Fresh bark samples of *F. religiosa* along with samples of *Panchavalka* tree were collected from various parts of India. DNA was isolated and purified using column. The bands that were found only in *F. religiosa* species and not in others were selected for further processing. The unique band thus obtained was further purified from the gel and used for cloning into T vector. The white colonies obtained were inoculated in LB media with antibiotic and plasmid isolation was done. The plasmids with retarded mobility were selected for confirmation with PCR using M13 primers. Positive clone was further purified and sequenced. The sequence obtained was used for designing SCAR primers. **Results:** For identification of *F. religiosa* by DNA marker DNA marker has been developed for *F. religiosa*. **Conclusion:** SCAR marker developed can be used to identify and differentiate the plant wherever it is used in preparation of medicines.

KEYWORDS *Ficus religiosa*, DNA SCAR marker, CTAB method, RAPD.

PICTORIAL ABSTRACT



ARTICLE HISTORY Received 28.11.2017 Accepted 30.04.2018

CORRESPONDENCE Dr Suma V Mallya, Associate professor, Department of Dravyaguna, SDM College of Ayurveda, Kuthpady, Udupi 574118, India. Email: sumamallya@gmail.com

CITE THIS RESEARCH AS Mallya SV, Udupi V, Prabhu SN, Sunil Kumar KN. Identification of *Ficus religiosa* By DNA SCAR Marker. J Ayu Med Sci 2018;3(1):340-3.

DOI 10.5530/jams.2018.3.5

Ficus religiosa Linn. is a large glabrous tree, without aerial roots from the branches, with sessile figs in axillary pairs, found all over India as an avenue tree^[1]. It is held sacred by Hindus and Buddhists. It belongs to family *Moraceae* and the genus *Ficus*, which is a large genus of trees or shrubs and about 65 species are reported from India^[2].

Bark of *F. religiosa* is popularly used in Indian system of medicine, for the treatment of many disease conditions. It is said to be astringent, antiseptic, laxative, haemostatic, vaginal disinfectant and used in diabetes, diarrhoea, leucorrhoea, menorrhagia, nervous diseases and also in skin problems^[3]. Bark contains beta-sitosterol, 1-D-glucosid, Vit K, n-otacosanol, methyl oleanolate, lanosterol, stigmasterol, lupen-3-one^[4]. Paste of the powdered bark is used as absorbent in inflammatory swellings. In Ceylon the juice of the bark is used as a mouth wash for toothache and for strengthening the gums. Water in which the freshly burnt ashes of the bark have been steeped is said to cure, obstinate cases of hiccup and stop vomiting sensation^[5]. Milk boiled with dried bark is a good aphrodisiac. Bark is also used as external application in many ailments. Bark powder is sprinkled over unhealthy ulcers and wounds to promote granulations. Rubbed with honey the powder is applied to apthous sores of children^[6]. *Nyagrodhadichoorna*, *Sarivadyasava*, *Karanjadighrita*, *Varadiguggulu* are few formulations prepared out of this drug^[7]. Bark occurs in flat or slightly curved pieces, varying from 1-2.5 cm or more in thickness. Outer surface brown or ash coloured surface uneven due to exfoliation of cork, inner surface smooth and somewhat brownish and fracture fibre^[8].

Bark drugs, because of similar macro-microscopic features are deliberately or accidentally admixed with other bark pieces of same species or other species^[9] most of the times. But each plant is combination of several constituents is having its own therapeutic property in treatment. Hence many pharmacognostic studies,

macro-microscopic atlas have been conducted and published in this regard.

Identification and quality assurance of botanical material is an essential requirement in herbal medicine to ensure reproducible quality of herbal medicine, which contributes to its safety and efficacy^[10]. Most of pharmacopoeia's, quality control guidelines suggest macroscopic, microscopic evaluation, chemical profiling for herbal drug standardization. Correct chemotype of plant can provide clinical efficacy. DNA markers are reliable in this regard as the genetic composition is unique for each species and is not affected by age, habitat, environmental factors and physiological conditions^[11]. Various types of DNA based molecular techniques are utilized to evaluate DNA polymorphism such as, DNA based molecular techniques, hybridization based methods, PCR based methods and sequencing based methods^[12].

Random Amplified Polymorphic DNA (RAPD) markers are DNA fragments from polymerase chain reaction amplification of random segments of genomic DNA with single primer of arbitrary nucleotide sequence^[13]. The limitation of RAPD is that it is not possible to distinguish whether a DNA segment is amplified from a locus that is heterozygous or homozygous. To overcome these problems SCAR(sequence characterized amplified region) marker is developed by cloning and sequencing RAPD marker(unique band) and further designing specific primers for the sequence obtained. These SCAR primers are used to identify and differentiate the strain of interest from others^[14].

Aim of present work is to develop SCAR primers for *Ficus religiosa* that is used in preparation of *Panchavalka* along with *F. benghalensis*, *F. glomerata*, *F. lacor* and *Thespesia populnea* bark powders. This would help in quality control stage of the formulations prepared from this important medicinal plant.

1. Plant material

Fresh bark samples of *F. religiosa* tree were collected from various parts of India. Plant samples were authenticated and Bark samples were frozen at -70°C till further use.

2. Preparation of bark powder

Bark samples were crushed into small pieces using autoclaved pestle and mortar and powdered thoroughly using dry ice intermittently. Individual pestle and mortar were used for each sample and collected the powder in a fresh container till further use.

3. DNA isolation

DNA isolation was attempted with different methods including urea method, DNAzol method, CTAB method and modified CTAB method. CTAB method DNA isolation was found better for RAPD PCR, though the yields were better with other methods. The crushed powder was treated with 70% ethanol before DNA extraction. Water wash to the powder was given to remove soluble pigments and compounds that may interfere in the experiment^[15].

4. CTAB method DNA extractions

Ethanol and water washed bark powder was homogenized with CTAB buffer thoroughly. Beta Mercapto ethanol was added and incubated for 90 minutes at 60°C in a water bath. This was centrifuged at 10,000 rpm for 15 minutes and supernatant was decanted into fresh tubes. Equal volume of 100% chloroform was added and mixed well for 10 minutes by inverting tube. This was, centrifuged at 10,000 rpm for five minutes. Separated aqueous layer was

transferred in to fresh tube and added equal volumes of iso-propyl alcohol. Centrifuged at 10,000 rpm for 15 minutes. Discarded supernatant and washed DNA pellets with 0.5 ml of 70% ethanol and centrifuged at 10,000 rpm for 5 minutes. Discarded ethanol and air dried DNA pellet. DNA was suspended in 1X TE buffer. The DNA was further purified using silica membrane based column and quantitated on agarosegel^[16].


5. RAPD PCR

Templates of all the 5 species were used for RAPD. PCR was set up using master mix consisting of 100µM each of dNTPs, 100mM KCl, 1.5mM MgCl₂, 0.1% each of tween 20 and Nonidet P40, 15p moles of random primer and 1u of taq polymerase in 40 µl volume. 100ng of template was used. PCR Cycle conditions were ,initial denaturation at 94°C for 5 minutes, denaturation for 30 seconds, annealing for 1 minute at 45°C, extension for 90 seconds at 72°C and final extension for another 7 minutes.

After amplification, the samples were run in 2% agarose gel using DNA marker with 0.1, 0.2, 0.3, 0.6, 1.0, 1.5, 2, 2.5, 3 and 3.5 Kb fragments.

Unique band of ~900bp was observed for *F. religiosa* which was not seen in other species. This fragment was gel purified , amplified and used for cloning into T vector. Clones were confirmed by amplification using vector primers and was sequenced. Sequence obtained was analysed and primers were designed to give ~550bp product for this region. Randomly selected ficus samples were used for checking the efficacy of the primers designed^[17].

Figure 1. RAPD with *Ficus* and *Thespesia* species



Lane1-4 racemosa, lane6-9. religiosa, lane11-14 benghalensis, lane15-17 thespacia, and lane18-20 lacor. Unique band is marked. M-DNA marker(0.1,0.2,0.3,0.6,1.0,1.5,2.0,2.5,3.0 and 3.5kb)

Samples were used from all the five *Panchavalka* plant species for RAPD (Figure 1). The bands that were found only in religiosa species and not in others were selected for further processing. The unique band thus obtained was further purified from the gel and used for cloning into T vector. The white colonies obtained were inoculated in LB media with antibiotic and plasmid isolation was done. The plasmids with retarded mobility were selected for confirmation with PCR using M13 primers. Positive clone was further purified and sequenced. The sequence obtained was used for designing SCAR primers. Highlighted sequence and primer designed is shown in Figure 2 and Table 1.

The primer selectively amplifies only *Ficus religiosa* samples (Figure 3). Thus SCAR marker for *F. religiosa* is successfully developed and can be used to identify and differentiate the same wherever it is used.

Figure 2. Clone confirmation 1,6 and 8 are clones




Fig: F. religiosa clones confirmed by vector primers

Table 1. The primers designed

ReFP	5'GAGCGAGAGAGGTAATGGAC'3
ReRP	5'CCATTGCTCGCAGCATTCT'3

6. Sequence data

ACGTCGCATGCTCCCGGCCGCTTCCCGTTCCCGTTCCTGAGAGCGAGAGA
GGTAATGGACGGCGGACGCGGCTTACCGCGGCGATGCCCTTCGGC
TTCTCCTCCTCTCTGCCACGTGCTCCTCTCAGGAGGAACGCCTGCG
ACGTCGCCCTTCGCCCCGAACGCTACAACCCGAAAACGACGATCGT
CGTCCGAATCCACTGCCGAGATACCGGTGCGCTCCACGGGATCATCATG
GTGTCGCCACCGGAGACCGAGGAAGGCGATGATAACAGCCACTCCCTCGA
GGTTTTTACACCCCGCCGGAGGAGTCGGTGGCTCAGAGCTCCAACGGG
CTGCAAGCGACTGTGGACGCGGTGACGGCGATGACGTTTGTGTCGGA
GTCCTGACGGCGGTGGTGAACGGTGGTTCGTTGGAGTTGGGAGCTCGGA
AGCCGATGGTGTGGGACGGTGGATTTGAGCCGGGATTCCGACCTAGGG
TTTCTGGAGGTTTCAGCCGACGACGAGGGAAAAAGACTATTTGGACTTGAA
TCGCTCGCTGTGGAGTTCCGGTGACCGAATCGAAGGGGGCTAGGGTT
TTGAGGAGGAATTGTCTTTGGACGACGGTTTGGGTAAGTACCATTGAA
GAAATTGAAGATTTAGAGCAAAATTTGGAAGAATCTGATGAAGAATTGG
ATCCTGAGAGAGCTACTCCTCCAAATCCCGGCTCTGCTCTCGCAAAATCTG
GTGAACAAAGCTCGGCGAAGAGGAAGTTGGAGTTCACCCAGAACGATAAT
TCCACCGAAGTCATCGAATGGGAATGAACAGACGAGGACGTTTTTGGT
GATGAGAAATGCTCGGAGCAATGAATAAATGGAGGCAACGGGATCCATG
GCCGCGGATATCACTAGTGCAGGCGCCCTGCAGGTGACCATATGGGAG
AGCTCCCAACGCGTTGGATGCATAG

Highlighted sequence is vector sequence. Red letters are flanking Nco I sites and rest of the sequence is of the insert that is sequence characterized for *Ficus religiosa*. Primer sequence is with green letters.

Quality control parameters with modern scientific techniques are of prime importance. The routine methods of herbal drug standardization address pharmacognostical, chemical, biological, biopharmaceutical and molecular approaches^[18].

Isolation of purified DNA from plant bark is challenging because of secondary metabolites and other compounds, which interfere with isolation process and purity^[19]. In this study DNA isolation was attempted with different methods and among these, CTAB method DNA isolation was found better for RAPD PCR, though the yields were better with other methods. The DNA obtained was further purified using silica membrane based column.

RAPD- a PCR based reaction introduced by Williams et al., which amplify segments of DNA those are essentially unknown to the researcher^[20]. Still the fragment polymorphisms produced by RAPD-PCR amplifications are not always reproducible. This limitation can be overcome by converting RAPD's into sequence characterized amplified region (SCAR)^[21].

In this study, templates of all the 5 species were used for RAPD. After amplification, the samples were run in 2% agarose gel using DNA marker with 0.1, 0.2, 0.3, 0.6, 1.0, 1.5, 2, 2.5, 3 and 3.5 Kb fragments. Unique band of ~900bp was observed for *F. religiosa* which was not seen in other species and this fragment was gel purified, amplified and used for cloning into T vector. Clones obtained were sequenced and primers were designed. The SCAR primers thus obtained were further validated using various ficus samples.

Figure 3. Amplification using SCAR primers



Fig: Ficus religiosa primer PCR

(Lane1: 500bp ladder, lane2-5 racemosa, lane6-9religiosa, 10-13 benghalensis, 14-16 lacor and 17-19 thespica.)

Clarity and reliability of SCAR markers proves them best techniques in current herbal drug standardization. SCAR primes of *Ficus religiosa* can be used to authenticate the sample, preventing its admixture with other bark drugs.

Acknowledgement Authors are grateful to Dr. D Veerendra Heggade, Revered President and Dr. Yashoverma B, Secretary, SDM Educational Society, Ujire for encouragement. Authors are grateful to Dr. Shreekanth U, Principal SDM college of Ayurveda Udupi and Dr. B Ravishankar, Director, SDM Centre for research in Ayurveda and Allied Sciences, Udupi for support and guidance. Authors are appreciative to Dr. Vasudha and Dr. Sudha Aristogene Biosciences Bangalore to carry out this research work.

Source of support Part of the observations is made from findings of a major research project entitled "Development of Sequence Characterized Amplified Region (SCAR) markers for the identification of popularly used five bark drugs 'Panchavalka', RGU:R&D:Res.Wing:2014-15 dated 13 MAR 2015, sanctioned by Rajiv Gandhi University of Health Sciences, Bangalore.

Conflict of interest Authors declare no conflict of interest

Contributors Dr. Suma Venkatesh Mallya collected plant samples and carried out DNA extraction, RAPD and SCAR marker at Aristogene Bangalore along with Dr.Vishwanatha. Dr.KN Sunil Kumar edited the manuscript giving suitable suggestions, to make it acceptable form. Suchitra Prabhu helped in authentication of plant samples.

References

1. Singh DD, Singh B, Goel RK. Traditional uses, phytochemistry and pharmacology of *Ficus religiosa*: A review. Journal of Ethnopharmacology 2011;565-583
2. Bhat KG. Flora of Udupi. Udupi: Indian Naturalist; 2003; p.597.
3. Nadakarni KM. Indian Materia Medica, Vol-I. Mumbai: Popular Prakashan Pvt Ltd, 1976; p.552.
4. Khare CP. Indian Medicinal Plants, An illustrated dictionary. New Delhi: Springer, 2007;p.265-9.
5. Kirtikar KR, Basu BD. Indian Medicinal Plants, Vol III. Dehradun: International Book Distributors; 2008; p.2317-8.
6. Bagade RJ, Bangale SV. Pharmacological studies in *Ficus religiosa*; IJPRS 2015;215-22.
7. K Nishteshwar. Text Book of Dravyaguna. Varanasi:Chaukamabha Surabharati Prakashana; 2007; p.541.

8. Data base on Medicinal Plants Used in Ayurveda, Vol III. New Delhi: CCRAS; 2005; p.131.
9. Gokhale SB, Kokate CH, Purohit AP. A text book of Pharmacognosy, 34th edition. Pune: Nirali Prakashan; 2013; p.4.9-4.11.
10. Quality control methods for medicinal plants. Geneva: World Health Organization; 1998; p.25-8.
11. Santoshkumar P, Ketkar P, Nayak S, Roy S. Application of DNA fingerprinting tools for authentication of Ayurvedic herbal medicine- A review. Journal of Scientific and Innovative Research 2014;606-12.
12. Raju BS, Manoj K, Sahoo D. Assessment of Genetic diversity of Ficus Species Using Rapd Markers as a Measure of Genomic Polymorphism; IOSR Journal of Pharmacy and Biological Sciences, 2015;22-7.
13. Sunil KumarKN, Maruthi KR, Alfarhan AH, Rajeshkrishnan R, Thomas J. Molecular fingerprinting of *Helianthus elastic* (Desr.) Danser growing on five different hosts by RAPD. Saudi Journal of Biological Sciences 2016;335-40.
14. Dhawan C, Kharb P, Sharma R, Uppal S, Aggraval RK. Development of male-specific SCAR marker in datepalm (*Phoenix dactylifera* L.). Tree Genetics and Genomics, 2013;1143-50.
15. Pirttila AM, Hirshikorpi M, KamarainenT, Jakkola L, Hohtola A. DNA isolation methods for medicinal and aromatic plants; Plant Molecular Biology Reporter2001;273a-f.
16. Sarvat M, Negi MS, Lakshmikumaran M, Tyagi AK, Das S, Srivastava PS. A standardized protocol for genomic DNA isolation from *Terminalia arjuna* for genetic diversity analysis. Electronic Journal of Biotechnology 2006;87-91.
17. Devaiah KM, Venkatasubramanian P. Genetic characterization and authentication of *Embelia ribes* using RAPD-PCR and SCAR marker. Planta Med 2008;1306-8.
18. Yadav NP, Dixit VK. Recent approaches in Herbal drug standardization. International Journal of Integrative Biology 2008;195.
19. Scot KD, Playford J. DNA extraction technique for PCR in rain forest plant species; Biotechniques 1996;974-8.
20. Joshi K, Chavan P, Warude D, Patwardhan B. Molecular markers in herbal drug technology. Current Science, 2004;159.
21. Polashock JJ. Development of SCAR markers for DNA fingerprinting and Germplasm analysis of American Cranberry. J AmerSocHortSci2002;677-84.